

PRACTICAL CLINICAL COURSES
*A Service of the Gordon J. Christensen
Career Development Program*

V3186
**Opening Vertical Dimension of Occlusion
(and Adjunctive Occlusal Procedures)**

Gordon J. Christensen, DDS, MSD, PhD

Materials Included:
C.E. Instruction Sheet
Products List
Clinician Responsible
Goals & Objectives
Overview
Supplemental Materials
AGD Post-Test

Gordon J. Christensen
PRACTICAL CLINICAL COURSES

**PROCEDURE FOR RECEIVING
CE CREDIT FOR VIDEO COURSES**

1. Complete the enclosed Post-Test.* For each **CE Video Purchased**, one complimentary test is included. Additional tests are \$35/person per test. Fees can be paid by check or with a credit card when submitting tests to Practical Clinical Courses. If more than 10 CE tests are required per video, please contact us at 800-223-6569.
2. Complete the demographic information located at the end of the test.
Type of Credit:
 - a. If the applicant selects "State," PCC will issue a certificate of verification to the applicant. The applicant must then submit this certificate to their state board to obtain credit.
 - b. If the applicant selects "AGD," PCC will submit credit information to the Academy of General Dentistry and confirmation to the applicant that it has been submitted (the applicant may check their AGD transcript for verification – please allow 30 days).
 - c. If the applicant selects "Both," PCC will complete a. & b. above.
3. Return the **Post-Test portion** via mail, fax, or email. Our contact information is as follows:

Practical Clinical Courses
3707 N Canyon Road
Suite 3D
Provo, UT 84604
Fax: (801) 226-8637
cetests@pccdental.com

4. Practical Clinical Courses will correct the Post-Test. **Passing scores are 70% or higher.**

****TO OBTAIN CE CREDIT ONLINE:*** Login or create an account on www.pccdental.com and select "My CE Tests" from the left-side menu. Click on the video title to take the test online. **RESULTS ARE IMMEDIATE.** Missing the test? Contact us at 800-223-6569 during our business hours of 7:00 a.m. – 4:00 p.m. MST to add it to your account.

Gordon J. Christensen
PRACTICAL CLINICAL COURSES
Sources of Products Discussed in
**V3186 Opening Vertical Dimension of Occlusion
(and Adjunctive Occlusal Procedures)**

Presented by: Gordon J. Christensen, DDS, MSD, PhD

1. **Access Blue**
Centrix, Inc.
770 River Road
Shelton, CT 06484
(800)235-5862
(203)929-5582
www.centrixdental.com
2. **AccuFilm II**
Parkell, Inc.
300 Executive Drive
Edgewood, NY 11717
(800)243-7446
(631)249-1134
www.parkell.com
3. **Blu-Mousse**
Parkell, Inc.
300 Executive Drive
Edgewood, NY 11717
(800)243-7446
(631)249-1134
www.parkell.com
4. **Caliper**
Local Dealer or
Architect Supplier
5. **Laboratory Occlusal Splints**
Various Laboratories
6. **Madame Butterfly Silk**
Almore International
441 19th Street S.E.
Hickory, NC 28602
(800)547-1511
www.almore.com
7. **NTI-tss 2.0**
Chairside Splint
(877)468-4877
www.chairsidesplintstore.com
8. **Occlusal Indicator Wax**
Kerr Corporation
200 S. Kraemer Blvd.
Building E2
Brea, CA 92821
(800)537-7123
www.kerrdental.com
9. **OptraGate**
Ivoclar Vivadent Inc.
175 Pineview Drive
Amherst, NY 14228
(800)533-6825
(716)691-0010
www.ivoclar.com
10. **QuickSplint**
Orofacial Therapeutics, Inc.
6824 El Cajon Blvd.
San Diego, CA 92115
(800)760-0526
www.quicksplint.com
11. **Megatray Custom Tray
Material (Replacement for
Triad TruTray)**
Henry Schein Dental
135 Duryea Road
Melville, NY 11747
(800)372-4346
(631)843-5500
www.henryschein.com
12. **Shimstock**
Almore International
441 19th Street S.E.
Hickory, NC 28602
(800)547-1511
www.almore.com
13. **Sideless Triple Tray**
Premier Dental Co.
1710 Romano Drive
Plymouth Meeting, PA 19462
(888)670-6100
(610)239-6000
www.premierdentalco.com
14. **Triad TruTray (See Megatray
Custom Tray Material)**
15. **TrollFoil by TrollDental**
Directa USA
64 Barnabas Road
Unit 3
Newtown, CT 06470
(800)537-8765
www.directausa.com

16. **T-Scan**
Tekscan, Inc.
333 Providence Hwy
Norwood, MA 02062
(800)248-3669
(617)464-4500
www.tekscan.com

Product names, the products themselves, and company names change rapidly. Please contact the companies shown to confirm current information.

Gordon J. Christensen Practical Clinical Courses, 3707 North Canyon Road, Suite 3D, Provo, UT 84604
Toll Free (800) 223-6569 or Utah Residents (801) 226-6569

The techniques and procedures on this videotape are intended to be suggestions only. Any licensed practitioner viewing this presentation must make his or her own professional decisions about specific treatment for patients. PCC is not responsible for any damages or other liabilities (including attorney's fees) resulting, or claimed to result in whole or in part, from actual or alleged problems arising out of the use of this presentation.

PROGRAM

V3186 Opening Vertical Dimension of Occlusion (and Adjunctive Occlusal Procedures)

CLINICIAN RESPONSIBLE:

Gordon J. Christensen, DDS, MSD, PhD
Founder and CEO, Practical Clinical Courses
Senior Consultant & Previous CEO, CR Foundation
Practicing Prosthodontist, Provo, Utah

GOALS & OBJECTIVES

At the completion of this video presentation, viewers should be able to:

1. List and discuss the current clinical terminology that defines occlusal positions and movements.
2. Define "normal" occlusion.
3. List 6 pathologic conditions of occlusion and their potential influence on vertical dimension of occlusion (VDO).
4. List the major factors that are known to reduce VDO.
5. Discuss grinding occlusion and its effect on VDO.
6. Discuss clenching bruxism and its effect on VDO.
7. Discuss gastroesophageal reflux and its effect on VDO.
8. Discuss diet and how it influences VDO in some cases.
9. Relate the wear resistance of restorative materials and the effect on VDO.
10. Discuss how periodontal disease can negatively influence VDO.
11. Discuss multiple tooth extractions and loss of VDO.
12. Describe the technique for locating VDO for an edentulous patient.
13. Describe the technique for locating VDO for a dentulous patient.
14. Describe speech tests and muscle tests for approximating VDO.
15. Define TENS and its use in locating VDO.
16. Discuss the importance of occlusal splints in locating VDO.
17. Discuss methods to transfer VDO from the patient to working casts.
18. List ten factors that influence the mesial-distal of tooth position.
19. List potential problems if VDO is opened too far.
20. List potential problems if VDO is not opened far enough.

OVERVIEW

V3186 Opening Vertical Dimension of Occlusion (and Adjunctive Occlusal Procedures)

Locating proper vertical dimension of occlusion (VDO) when rehabilitating patients is a challenging task, and is often done incorrectly. Such improper positioning can cause serious patient discomfort, muscle dysfunction, and potential destruction of the rehabilitation.

The following and other topics are included in this presentation:

- Occlusal terminology
- What is normal occlusion
- Six pathologic conditions of occlusion and their influence on VDO
- Major factors that reduce VDO
- Grinding bruxism
- Clenching bruxism
- Diet and its influence on VDO
- Gastroesophageal reflux disease, and its effect on VDO
- Wear of restorative materials and its effect on VDO
- Periodontal disease and its effect on VDO
- Multiple tooth extractions and the effect on VDO
- Diagnostic aids for occlusal examination and treatment
- Opening vertical dimension of occlusion
- Finding rest position
- Speech tests that help to locate VDO
- The effect of TENS on finding VDO
- The usefulness of mounted diagnostic casts
- Occlusal splints and their usefulness in treatment of VDO determination
- Testing muscle activity for determining VDO
- Identification of reasons for collapsed VDO and future prevention
- Interocclusal records
- Methods to transfer VDO to working casts
- Factors altering mesial-distal dimension and their effect on VDO

SUPPLEMENTAL MATERIALS

V3186 Opening Vertical Dimension of Occlusion (and Adjunctive Occlusal Procedures)

1. Alhadj MN, Khalifa N, Abduo J, Amran AG, Ismail IA. Determination of occlusal vertical dimension for complete dentures patients: an updated review. *J Oral Rehabil.* 2017 Nov;44(11):896-907. doi: 10.1111/joor.12522. Epub 2017 Jun 10.
2. Remiszewski D, Bidra AS, Litt MD. Ability of General Dentists and Prosthodontists to Discern and Identify Incremental Increases in Occlusal Vertical Dimension in Dentate Subjects. *Int J Prosthodont.* 2017 Jul/Aug;30(4):327-33. doi: 10.11607/ijp.5152.
3. Hahnel S, Scherl C, Rosentritt M. Interim rehabilitation of occlusal vertical dimension using a double-crown-retained removable dental prosthesis with polyetheretherketone framework. *J Prosthet Dent.* 2017 Jun 20. pii: S0022-3913(17)30155-5. doi: 10.1016/j.prosdent.2017.02.017. [Epub ahead of print]
4. Ting J, Shuhui H, Hongqiang Y, Lu J. CAD/CAM Ceramic Overlays to Restore Reduced Vertical Dimension of Occlusion Resulting from Worn Dentitions: A Case History Report. *Int J Prosthodont.* 2017 May/Jun;30(3):238-41. doi: 10.11607/ijp.5146.
5. Alhadj MN, Daer AA. A proposed linear skeletal distance to predict occlusal vertical dimension: A cephalometric study. *J Prosthet Dent.* 2017 Apr 20. pii: S0022-3913(17)30059-8. doi: 10.1016/j.prosdent.2016.12.022. [Epub ahead of print]
6. Christensen GJ. Ask Dr. Christensen: WHAT CAUSES CHANGES IN OCCLUSION? *Dental Economics.* 2013 Jul;103(7):24-31.
7. Christensen GJ. Ask Dr. Christensen: TREATMENT OF CRACKED TEETH. *Dental Economics.* 2011 Apr;101(4):68-74.
8. Christensen GJ. The major part of dentistry you may be neglecting. *J Am Dent Assoc.* 2005 Apr;136(4):497-9.
9. Christensen GJ. Is occlusion becoming more confusing? A plea for simplicity. *J Am Dent Assoc.* 2004 Jun;135(6):767-70.

SUPPLEMENTAL MATERIALS (CONT'D)

**V3186 Opening Vertical Dimension of Occlusion
(and Adjunctive Occlusal Procedures)**

10. Christensen GJ. Now is the time to observe and treat dental occlusion. J Am Dent Assoc. 2001 Jan;132(1):100-2.
11. Christensen GJ. TREATING BRUXISM AND CLENCHING. J Am Dent Assoc. 2000 Feb;131(2):233-5.
12. Christensen GJ. ABNORMAL OCCLUSAL CONDITIONS: A FORGOTTEN PART OF DENTISTRY. JADA. 1995 Dec;126(12):1667-8.

POST-TEST

V3186 Opening Vertical Dimension of Occlusion (and Adjunctive Occlusal Procedures)

1. Reduced canine rise and reduced incisal guidance are present in:
 - a. secondary occlusal trauma.
 - b. grinding bruxism.
 - c. clenching bruxism.
 - d. primary occlusal trauma.

2. Which condition is associated with advanced periodontal disease?
 - a. Primary occlusal trauma
 - b. Grinding bruxism
 - c. Clenching bruxism
 - d. Secondary occlusal trauma

3. Rest position of the mandible is usually ____ mm from maximum intercuspation.
 - a. 0.5
 - b. 1.0
 - c. 1.5
 - d. None of the above

4. The sound "S" usually places the maxillary and mandibular teeth about _____ mm apart.
 - a. 5
 - b. 4
 - c. 1
 - d. None of the above

5. Which is incorrect relative to TENS (transcutaneous electrical nerve stimulation)?
 - a. Can assist in relaxing the muscles of mastication and finding VDO.
 - b. Requires about ½ hour of application to relax muscles.
 - c. Is not potentially harmful if used on a person with a pacemaker.
 - d. Does not require an expensive device.

6. Which is usually the most destructive condition relative to closing VDO?
 - a. Diet
 - b. Grinding bruxism
 - c. Primary occlusal trauma
 - d. Wear on ceramic restorations

7. When VDO is opened too much, the following occurs:
 - a. near total inactivation of both the masseter and temporalis muscles.
 - b. near total inactivation of the temporalis muscles.
 - c. near total inactivation of the masseter muscles.
 - d. none of the above.

POST-TEST (CONT'D)

**V3186 Opening Vertical Dimension of Occlusion
(and Adjunctive Occlusal Procedures)**

8. When VDO is closed too much, the following occurs:
 - a. near total inactivation of the temporalis muscles.
 - b. near total inactivation of the masseter muscles.
 - c. near total inactivation of both the masseter and temporalis muscles.
 - d. none of the above.

9. Which method is almost always necessary to determine the amount of opening of VDO that is tolerable to a patient?
 - a. Occlusal equilibration
 - b. TENS
 - c. Occlusal splint
 - d. Muscle manipulation

10. One of the easiest procedures to determine if occlusal contacts on a crown are too low or too high is:
 - a. mounted casts.
 - b. use of articulating papers.
 - c. spray-on articulating media.
 - d. TENS.
 - e. Kerr Occlusal Indicator Wax.

PLEASE PRINT

Name _____

Name of video purchaser (if different from above) _____

Address _____

City/State/Zip _____

Phone No. _____

Email _____

Indicate which type of credit you wish to obtain: AGD State Both

State License No. _____ AGD No. _____

Payment information Visa American Express Mastercard Discover

Card # _____ - _____ - _____ - _____ Expires ____/____/____ CVV2-Code: _____

For each **CE Video Purchased**, one complimentary test is included. If you require CE tests for *staff members or additional doctors to receive credit, the fee is \$35/person per test.*

If more than 10 CE tests are required per video, please contact PCC at 800-223-6569.